

8200006

THE UNITED SHAMES OF AMIERICAL

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Northrup King Company

Withereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eighteen years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to excide others from selling the variety, or offering it for sale, or reproducing it, porting it, or exporting it, or using it in producing a hybrid or different therefrom, to the extent provided by the Plant Variety Protection Act 1942, as amended, 7 u.s.c. 2321 et seq.)

COMMON WHEAT

18351

In Testimony Waterrot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington this 11th day of March in the year of our Lord one thousand nine hundred and eighty-two.

Allest:

Kerneth H. E.
Acting
Commissioner
Plant Variety Protection Office
Grain Division
Agricultural Marketing Service

John R Block Secretary of Agriculture

	UNITED STATES DEPARTME	CETING SERVICE			FORM APPROVED OMB NO. 40-R3822
	LIVESTOCK, POULTRY, GRA PLICATION FOR PLANT VARIE TRUCTIONS: See Reverse.			No certificate for pla be issued unless a co has been received (5 l	ant variety protection may impleted application form
-	TEMPORARY DESIGNATION OF VARIETY	1b. VARIETY NAM	E		AL USE ONLY
	75W 355	835		PV NUMBER	390006
2.	KIND NAME	3. GENUS AND SPE	CIES NAME	FILING DATE	TIME A.M.
	Common Wheat	Triticum aesti	<u>vum</u> Lin.≔	10/19/81 FEE RECEIVED	2:00 (P.M.) DATE
4.	FAMILY NAME (BOTANICAL) Gramineae	5. DATE OF DETE		\$ 500.00 \$ 250.00	10/19/81 1/11/82
6,	NAME OF APPLICANT(S)	· · · · · · · · · · · · · · · · · · ·	t and No. or R.F.D. No.,	City State and ZIP	8. TELEPHONE AREA
	Northrup King Co.	Code)	1500 Jackson St.	• •	CODE AND NUMBER
	in ap imig		Minneapolis, MN		612-781-5305
9.	IF THE NAMED APPLICANT IS NOT A PE ORGANIZATION: (Corporation, partnersh.	RSON, FORM OF ip, association, etc.)	10. IF INCORPORATI	ED, GIVE STATE AND PORATION	11. DATE OF INCOR- PORATION
	Corporation		Dela		1896
12.	N 1	lobert W. Romig Iorthrup King C 500 Jackson St.	5		ATION AND RECEIVE
13.	CHECK BOX BELOW FOR EACH ATTACH				
	13A. Exhibit A, Origin and Bree	ding History of the	Variety (See Section 5	12 of the Plant Variety	Protection Act.)
	X 13B. Exhibit B, Novelty Statem			_	·
	13C. Exhibit C, Objective Descri	intion of the Variets	Request form from	Dlant Variato Protect	ion Office)
				rumi vanety Protecti	ion Office.)
	13D. Exhibit D, Additional Desc	cription of the Varie	ty.		
14a.	DOES THE APPLICANT(S) SPECIFY THAT SEED? (See Section 83(a). (If "Yes," answer	r SEED OF THIS VAR er 14B and 14C below.)	IETY BE SOLD BY VAF		A CLASS OF CERTIFIED
14b.	DOES THE APPLICANT(S) SPECIFY THAT LIMITED AS TO NUMBER OF GENERATI	T THIS VARIETY BE	14c. IF "YES," TO 14I	B, HOW MANY GENERA	ATIONS OF PRODUC-
	YES NO	0.40.	FOUNDATION	REGISTERED	CERTIFIED
15a.	DID THE APPLICANT(S) FILE FOR PROTI	ECTION OF THIS VAR			NO (If "Yes," give
	name of countries and dates.)				B12(4/2)
15b.	HAVE RIGHTS BEEN GRANTED THIS VA and dates.)	RIETY IN OTHER CO	UNTRIES? YES	XNO (If "Yes,"	give name of countries
	•				
16.	DOES THE APPLICANT(S) AGREE TO THE JOURNAL? The applicant(s) de Leu (s) de la constant de la co				
17,	The applicant(s) declare(s) that a viable replenished upon request in accordance	with such regulation	ns as may be applicabl	le.	
. :	The undersigned applicant(s) is (are) the variety is distinct, uniform, and stable a 42 of the Plant Variety Act.	e owner(s) of this se is required in Section	xually reproduced nov 1 41, and is entitled to	vel plant variety, and loop protection under the	believe(s) that the provisions of Section
	Applicant(s) is (are) informed that false	representation here	in can jeop <i>ati</i> lize prot	ection and result in p	englties.
0	CTOBER IS. 1981	200	Hober	1-11- M	ones,
	(DATE)		(S	SIGNATURE OF APPLIC	CANT)
					_

EXHIBIT A Origin and Breeding History of the Variety

Variety 835 is the result of hybridization, individual plant selection and head selection from the cross Lancer/Sturdy. Our pedigree for this variety is N7015-2K-OK.

We made the cross in the greenhouse at Eden Prairie, Minnesota, in 1971 and grew the F1 in southern California during 1971 by vernalizing the seed and transplanting F1 seedlings in the field. F2 seed was planted at Pratt, Kansas in the fall of 1972. Individual F2 plant selection was made in 1973. The F3 plant row was bulked at Pratt in 1974 and the F4 was tested in 1975 preliminary yield trials. The F5 seed from the preliminary trial was utilized to plant replicated trials in 1975-76. In 1976 twenty heads were selected from the replicated yield trial (F5) at Pratt, Kansas. The heads were threshed individually and F6 head-rows were grown at Yuma in 1976-Ten head-row lines were harvested at Yuma, Arizona. Seed from these individual head-row lines was then yield tested in replicated trials at Pratt, Kansas and York, Nebraska in 1977-78. An increase of each of the ten head-row lots was simultaneously grown at Yuma in 1977-78. Four of the head-row lines were tested again in 1978-79 and increased at Yuma as pure lines. In August, 1979, one head-row line, 79AWH 3003, was selected to represent the variety, based on yield and agronomic data. 835 is thus derived from a single F6 head-row selection. Breeder seed in 1980 was in the F9 generation of selfing.

The variety is uniform and stable. There are no unusual or characteristic variations.

Foundation seed produced in 1980 has been inspected and approved by the Texas Seed Certification Program.

EXHIBIT B Novelty Statement

Variety 835 is most similar to Scout 66 but differs in plant height. Variety 835 is a short stature wheat with a plant height averaging 81-91 cm depending upon environment. The height of Scout 66 ranges from 106-110 cm in the same environments. Variety 835 is thus from 19-25 cm shorter than Scout 66.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, POULTRY, GRAIN & SEED DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C (Wheat)

OBJECTIVE DESCRIPTION OF VARIETY

INSTRUCTIONS: See Nevelsel	ITICUM SPP.)
NAME OF APPLICANT(S)	FOR OFFICIAL USE ONLY
Northrup King Co., ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	PVPO NUMBER
	VARIETY NAME OR TEMPORARY
Minneapolis MN 55413	DESIGNATION
	835
Place the appropriate number that describes the varietal character. Place a zero in first box (e.g. 0 8 9 or 0 9) when number	
1. KIND:	
1 1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5	= POLISH 6 = POULARD 7 = CLUB
2. TYPE;	1 = SOFT 3 = OTHER (Specify)
2 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	
2 1 = WHITE 2 = RED 3 = OTHER (Specify)	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
FIRST FLOWERING	LAST FLOWERING
4. MATURITY (50% Flowering):	
0 1 NO. OF DAYS EARLIER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS
0 1 NO. OF DAYS LATER THAN	4 = LEMHI 5 = NUGAINES 6 = LEEDS 7=Centurk 8=Vona
5. PLANT HEIGHT (From soil level to top of head):	
0 9 1 CM. HIGH	
0 0 CM. TALLER THAN	8 7=Centurk 8=Vona
CM. SHORTER THAN	
6. PLANT COLOR AT BOOTING (See reverse):	7. ANTHER COLOR:
2 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 1 = YELLOW 2 = PURPLE
8. STEM:	
1 Anthocyanin: l = ABSENT 2 = PRESENT	2 Waxy bloom: 1 = ABSENT 2 = PRESENT
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT	1 Internodes: 1 = HOLLOW 2 = SOLID
0 3 NO. OF NODES (Originating from node above ground)	2 0 CM, INTERNODE LENGTH BETWEEN FLAG LEAF
9. AURICLES:	
Anthocyanin: = ABSENT 2 = PRESENT	2 Hairiness: 1 = ABSENT 2 = PRESENT
O. LEAF:	
Flag leaf at = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):	Flag leaf: 1 = NOT TWISTED 2 = TWISTED
Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT	2 Waxy bloom of flag leaf sheath: ABSENT 2 = PRESENT
0 9 MM. LEAF WIDTH (First leaf below flag leaf)	2 2 CM. LEAF LENGTH (First leaf below flag leaf):

11. HEAD:	•	•	
Density: 1 = LAX	2 = DENSE	Shape: 1 = TAPER 4 = OTHER	ING 2 = STRAP 3 = CLAVATE (Specify)
4 Awnedness: 1 = Av	NLESS 2 = APICALLY AWNLETED	3 = AWNLETED 4 = AWNE	D
Color at maturity: 5	= WHITE 2 = YELLOW 3 = PINK 4 = BROWN 6 = BLACK 7 = OTHE	= RED ER (Specify):	
0 7 CM. LENGTH		0 7 MM. WIDTH	
12. GLUMES AT MATUR Length: 1 = SHORT 3 = LONG		2 Width: 1 = NARRO 3 = WIDE (C	
Shoulder 1 = WANT shape: 4 = SQUA			2 = ACUTE 3 = ACUMINATE
13. COLEOPTILE COLOR	l:	14. SEEDLING ANTHOCY	'ANIN:
1 = WHITE 2 = R	ED 3 = PURPLE		= PRESENT
15. JUVENILE PLANT GR	OWTH HABIT:	<u> </u>	
1 = PROSTRATE	2 = SEMI-ERECT 3 = EREC	et .	
16. SEED:			
1 Shape: I = OVATE	2 = OVAL 3 = ELLIPTICAL	1 Cheek: I = ROUND	ED 2 = ANGULAR
2 Brush: 1 = SHORT	2 = MEDIUM 3 = LONG	Brush: 1 = NOT CO	DLLARED 2 = COLLARED
Phenol reaction (See instructions):	1 = IVORY 2 = FAWN 3 = LT. BROWN 4 = BROWN 5 = BLACK	N	
3 Color: I = WHITE	2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specify)	
0 6 MM. LENGTH	0 3 MM. WIDTH	3 6 GM. PER 1000	SEEDS
17. SEED CREASE:			
	ESS OF KERNEL 'WINOKA'	Depth: 1 = 20% Of	R LESS OF KERNEL 'SCOUT'
2 = 80% OR LE	ESS OF KERNEL 'CHRIS'		LESS OF KERNEL 'CHRIS'
	AS WIDE AS KERNEL 'LEMH!'	3 = 50% OF	LESS OF KERNEL 'LEMHI'
I8. DISEASE: (0 = Not Test	ed, 1 = Susceptible, 2 = Resistant)		
1 (Races) Seedling test	OSH LEAF RUST Unknown	STRIPE RUST	0 LOOSE SMUT
0 POWDERY MILDEW	0 BUNT	1 OTHER (Specify) S	oil-borne mosaic virus
19. INSECT: (0 = Not Teste	d, 1 = Susceptible, 2 = Resistant)		
O SAWFLY	0 APHID (Bydv.)	O GREEN BUG	O CEREAL LEAF BEETLE
OTHER (Specify)	HESSIAN FLY	1 GP 0 A	0 в 0 с
	RACES:		
•	,	0 D E	0 F 0 G
20. INDICATE WHICH VARIE	TY MOST CLOSELY RESEMBLES THAT S	UBMITTED:	
CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant tillering	Scout	Seed size	Scout
Leaf size	Centurk	Seed shape	Scout
Leaf color	Sturdy	Coleoptile elongation	Scout
Leaf carriage	Sturdy	Seedling pigmentation	Scout

INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

LEAF COLOR: Nickerson's or any recognized color fan should be used to determine the leaf color of the described variety.

EXHIBIT D Additional Description of the Variety

Variety 835 is a cultivar of <u>Triticum aestivum</u> L. with winter growth habit. The kernels are hard, red, and ovate to elliptical in shape. Cheeks are rounded normally. The brush is midsized and midlong. Spikes are awned, lax to middense, and fusiform to oblong. Glumes are white and glabrous. The beak is accuminate and short (1-4 mm long).

This variety is demidwarf with height similar to Vona. Date of heading for 835 averages one day earlier than Centurk and one day later than Vona. Relative maturity is medium. 835 has the adult plant resistance of Sturdy to leaf rust (Puccinia recondita) in the field. Adult plant resistance to some races of stem rust (P. graminis f. sp. tritici) has been observed in the field. However seedling tests to races 15B2 and 151QSH indicate susceptible reactions (rating of 3). 835 is moderately susceptible to the Great Plains Biotype of Hessian Fly (Mayetiola destructor (Say)).

Coleoptile color is white and seedling anthocyanin is absent. Plant color at booting is green. The stem is hollow and has no anthocyanin. Usually three nodes are formed above the ground.

Overall quality for bread has been comparable to check varieties. Stability and valorimeter scores have often been greater for 835 than for Centurk.

835 is adapted to the hard red winter wheat growing areas of Texas, Oklahoma, Kansas, Colorado, Nebraska, and New Mexico.

Table 1. Plant Height Comparisons of Variety 835 with Check Varieties in 1980 Nebraska Variety Tests

	Plant Height (cm)					
Location	835	Scout 66	Ctk 78	Bennett	Vona	
Clay Co.	89	114	122	107	94	
Kearney Co.	76	102	91	89	76	
Lincoln Co.	84	102	99	91	86	
Hayes Co.	89	117	112	104	94	
Keith Co.	<u>69</u>	94	<u>86</u>	74	<u>69</u>	
Average	81	106	102	93	84	

Table 2. Plant Height Comparisons of Variety 835 with Check Varieties in the 1980 Texas Hard Red Winter Wheat Test

		Plant Height (cm)				
Location	835	Tam 101	Newton	Vona		
Bushland Dry	51	61	58	56		
Bushland Ir.	76	76	76	74		
Chillicothe	58	56	61	58		
Munday	41	41	38	41		
Abiliene	<u>43</u>	48		<u>46</u>		
Average 5 Loc.	54	56	56	55		

Table 4. Agronomic Characteristics of 835 in Comparison to Checks Summarized from Northrup King Trials in 1978 and 1979

	835	Centurk	Vona	Scout 66
Height cm	91	105	91	110
Test Wt. kg/hl	74.5	74.0	71.7	74.1
Heading Date	144	146	143	145
Lodging (0-9) <u>1</u> /	2	4	2	6
Winter Survival (0-9) 2/	8	8	5	8
Shattering (0-9) <u>1</u> /	2	2	2	3
Leaf Rust (Field)	MR	s	MS	s
Stem Rust (Field)	R	R	R	R
Soil Borne Mosaic Virus	s	MS	S	S

^{1/0-9} scale where 0 is best and 9 is poorest

 $[\]underline{2}$ / 0-9 scale where 0 = no survival; 1 = 10-19% survival; 9 = 90-100% survival

Table 5. Quality Characteristics of Samples Grown at York, Nebraska, '78 and '79

	1978		1979	
Characteristics	835	Centurk	835	Centurk
Wheat Protein	16.05	14.0	13.80	13.75
Test Weight	52.6	58.1	63.2	62.7
Milling & Ext.	67.2 F	68.5 G-	71.2 G	69.0 G
Farinograph				
Absorption	63.0	61.0	57.0	60.4
Peak	25.0	11.0	8.0	6.25
Stability	40.0	29.0	26.0	16.5
MTI	10	10	20	25
Valorimeter	100	83	73	65
Flour				
Ash	.441	.388	.406	.352
Protein	14.75	12.85	12.80	12.60
Bake				
Absorption	65.5 G+	64.0 G	59.5 F	63.0 F
Mix	7.25	8.25 F	5.75 G-	5.00 V
Dough	5 G	5 G-	6 G	6 G
Loaf Vol. cc	1000 Ex-	995 G+	980 VG	1000 EX
Score	28 G-	29 G-	31 G-	35 VG-
Overall Score	57 G-	60 G-	60 G-	64 G

Table 6. Quality Characteristics of Samples Grown at Pratt Kansas in '78 and '79

835	Centurk
12.00	12.60
63.5	59.9
72.9 G	67.5 F
57.7	59.0
8.00	8.00
23.50	45.00
30	20
74.	75
.438	.379
11.00	11.35
	,
61.0 G-	62.5 G-
5.00 VG	6.50 G-
5 G-	5 G-
	900 G
30 G-	28 G-
60 G-	55 G-
	880 G- 30 G-